arXiv:1203.4899v2 [math.RT] 14 May 2012

The algebra of invariants of the adjoint action of the unitriangular group in the nilradical of a parabolic subalgebra

V. V. Sevostyanova*

Abstract. In the paper the algebra of invariants of the adjoint action of the unitriangular group in the nilradical of a parabolic subalgebra is studied. We set up a conjecture on the structure of the algebra of invariants. The conjecture is proved for parabolic subalgebras of special types.

Consider the general linear group $\mathrm{GL}(n,K)$ defined over an algebraically closed field K of characteristic 0. Let B (N, respectively) be its Borel (maximal unipotent, respectively) subgroup, which consists of triangular matrices with nonzero (unit, respectively) elements on the diagonal. We fix a parabolic subgroup P that contains B. Denote by $\mathfrak{p}$, $\mathfrak{b}$ and $\mathfrak{n}$ the Lie subalgebras in $\mathfrak{gl}(n,K)$ that correspond to P, B and N, respectively. We represent $\mathfrak{p}=\mathfrak{r}\oplus\mathfrak{m}$ as the direct sum of the nilradical $\mathfrak{m}$ and a block diagonal subalgebra $\mathfrak{r}$ with sizes of blocks $(n_1,\ldots,n_s)$. The subalgebra $\mathfrak{m}$ is invariant relative to the adjoint action of the group P, therefore $\mathfrak{m}$ is invariant relative to the adjoint action of the subgroups B and N. We extend this action to the representation in the algebra $K[\mathfrak{m}]$ and in the field $K(\mathfrak{m})$. The subalgebra $\mathfrak{m}$ contains a Zariski-open P-orbit, which is called the *Richardson orbit* (see [R]). Consequently, $K[\mathfrak{m}]^P=K$. In this paper, we study the structure of the algebra of invariants $K[\mathfrak{m}]^N$. In the case $P=B$, the algebra of invariants $K[\mathfrak{m}]^N$ is the polynomial algebra $K[x_{12},x_{23},\ldots,x_{n-1,n}]$. Let $\mathfrak{r}$ be the sum of two blocks; this case was considered by M. Brion in the paper [B]. A complete description of the field of invariants $K(\mathfrak{m})^N$ for any parabolic subalgebra is a result of [S]. The question concerning the structure of the algebra of invariants $K[\mathfrak{m}]^N$ remains open and seems to be a considerable challenge. We do not know when the algebra of invariants $K(\mathfrak{m})^N$ is finitely generated.

---

*This research was partially supported by the RFBR (project 08-01-00151-a)

In this paper, we consider the series of polynomials (see the notations (5), (6), (7)) in the algebra of invariants. We show that these polynomials generate the algebra of invariants in special cases (Theorems 4.1 and 5.2). We state a conjecture on the structure of the algebra of invariants $K[\mathfrak{m}]^N$ (Conjecture 5.4).

§1. The main definitions

We begin with definitions. Every positive root γ in $\mathfrak{gl}(n, K)$ has the form (see [GG]) $\gamma = \varepsilon_i - \varepsilon_j$, $1 \leqslant i < j \leqslant n$. We identity a root γ with the pair (i, j) and the set of the positive roots Δ^+ with the set of pairs (i, j), $i < j$. The system of positive roots $\Delta^+_{\mathfrak{r}}$ of the reductive subalgebra $\mathfrak{r}$ is a subsystem in Δ^+.

Let $\{E_{i,j} :\ i < j\}$ be the standard basis in $\mathfrak{n}$. By E_γ denote the basis element $E_{i,j}$, where $\gamma = (i, j)$.

We define a relation in Δ^+ such that $\gamma' \succ \gamma$ whenever $\gamma' - \gamma \in \Delta^+_{\mathfrak{r}}$. If $\gamma \prec \gamma'$ or $\gamma \succ \gamma'$, then the roots γ and γ' are said to be *comparable.* Denote by M the set of $\gamma \in \Delta^+$ such that $E_\gamma \in \mathfrak{m}$. We identify the algebra $K[\mathfrak{m}]$ with the polynomial algebra in the variables $x_{i,j},\ (i, j) \in M$.

Definition 1.1. A subset S in M is called a *base* if the elements in S are not pairwise comparable and for any $\gamma \in M \setminus S$ there exists $\xi \in S$ such that $\gamma \succ \xi$.

Definition 1.2. Let A be a subset in S. We say that γ is a *minimal element* in A if there is no $\xi \in A$ such that $\gamma \succ \xi$.

Note that M has a unique base S, which can be constructed in the following way. We form the set S_1 of minimal elements in M. By definition, $S_1 \subset S$. Then we form a set M_1, which is obtained from M by deleting S_1 and all elements

$$\{\gamma \in M : \exists\ \xi \in S_1,\ \gamma \succ \xi\}.$$

The set of minimal elements S_2 in M_1 is also contained in S, and so on. Continuing the process, we get the base S.

Definition 1.3. An ordered set of positive roots $\{\gamma_1, \ldots, \gamma_s\}$ is called a *chain* if $\gamma_1 = (a_1, a_2)$, $\gamma_2 = (a_2, a_3)$, $\gamma_3 = (a_3, a_4)$, and so on. The number s is called the *length of a chain.*

Definition 1.4. We say that two roots $\xi, \xi' \in S$ form an *admissible pair* $q = (\xi, \xi')$ if there exists $\alpha_q \in \Delta^+_{\mathfrak{r}}$ such that the ordered set of roots $\{\xi, \alpha_q, \xi'\}$ is a chain. Note that the root α_q is uniquely determined by q.

We form the set $Q := Q(\mathfrak{p})$ that consists of admissible pairs of roots in S. For every admissible pair $q = (\xi, \xi')$ we construct a positive root $\varphi_q = \alpha_q + \xi'$. Consider the subset $\Phi = \{\varphi_q : \ q \in Q\}$ in the set of positive roots.

Definition 1.5. The set $S \cup \Phi$ is called an *expanded base.*

Using the given parabolic subgroup, we construct a diagram, which is a square matrix in which the roots from S are marked by the symbol $\otimes$ and the roots from Φ are labeled by the symbol $\times$. The other entries in the diagram are empty.

Example 1.6. Below a diagram for a parabolic subalgebra with sizes of its diagonal blocks $(2,1,3,2)$ is given.



Diagram 1

Consider the formal matrix $\mathbb{X}$ in which the variables $x_{i,j}$ occupy the positions $(i,j) \in M$ and the other entries are equal to zero. For any root $\gamma = (a,b) \in M$ we denote by S_γ the set of $\xi = (i,j) \in S$ such that $i > a$ и $j < b$. Let $S_\gamma = \{(i_1, j_1), \ldots, (i_k, j_k)\}$. Denote by M_γ a minor M_I^J of the matrix $\mathbb{X}$ with ordered systems of rows I and columns J, where

$$I = \mathrm{ord}\{a, i_1, \ldots, i_k\}, \quad J = \mathrm{ord}\{j_1, \ldots, j_k, b\}.$$

For every admissible pair $q = (\xi, \xi')$ such that q corresponds to $\varphi \in \Phi$, we construct the polynomial

$$L_\varphi = \sum_{\substack{\alpha_1, \alpha_2 \in \Delta_{\mathfrak{r}}^+ \cup \{0\} \\ \alpha_1 + \alpha_2 = \alpha_q}} M_{\xi + \alpha_1} M_{\alpha_2 + \xi'}. \qquad (1)$$

Theorem 1.7. [PS] *For an arbitrary parabolic subalgebra, the system of polynomials*

$$\{M_\xi, \ \xi \in S, \ L_\varphi, \ \varphi \in \Phi, \}$$

is contained in $K[\mathfrak{m}]^N$ *and is algebraically independent over* K.

Denote by $\mathcal{Y}$ the subset in $\mathfrak{m}$ that consists of matrices of the form

$$\sum_{\xi\in S} c_\xi E_\xi + \sum_{\varphi\in\Phi} c'_\varphi E_\varphi,$$

where $c_\xi \neq 0$ and $c'_\varphi \neq 0$.

Definition 1.8. The matrices from $\mathcal{Y}$ are said to be *canonical.*

The proof of the following theorem is found in [S].

Theorem 1.9. *There exists a nonempty Zariski-open subset $U \subset \mathfrak{m}$ such that the N-orbit of any $x \in U$ intersects $\mathcal{Y}$ at a unique point.*

Now let $\mathcal{S}$ be the set of denominators generated by minors M_ξ, $\xi \in S$. Consider the localization $K[\mathfrak{m}]^N_{\mathcal{S}}$ of the algebra of invariants $K[\mathfrak{m}]^N$ with respect to $\mathcal{S}$. Since the minors M_ξ are N-invariants, we have

$$K[\mathfrak{m}]^N_{\mathcal{S}} = \big(K[\mathfrak{m}]_{\mathcal{S}}\big)^N.$$

The following results are consequences of Theorem 1.9.

Theorem 1.10. *The ring $K[\mathfrak{m}]^N_{\mathcal{S}}$ is the ring of polynomials in $M_\xi^{\pm1}$, $\xi \in S$, and in L_φ, $\varphi \in \Phi$.*

Theorem 1.11. *The field of invariants $K(\mathfrak{m})^N$ is the field of rational functions of M_ξ, $\xi \in S$, and L_φ, $\varphi \in \Phi$.*

The polynomials M_ξ and L_φ do not generate the algebra of invariants. We prove in §4 and §5, respectively, that if the reductive subalgebra $\mathfrak{r}$ consists of three blocks or of four blocks with sizes $(1,2,2,1)$, then the algebra of invariants $K[\mathfrak{m}]^N$ is not generated by the invariants M_ξ and L_φ. Let the reductive subalgebra consist of blocks with sizes $(2,k,2)$, $k > 3$, or with sizes $(1,2,2,1)$. We construct a polynomial D such that the algebra of invariants is generated by D and by the invariants M_ξ and L_φ.

§2. The other definition of the invariant L_φ

In the present chapter, we give the alternative definition of the invariant L_φ, where $\varphi \in \Phi$.

Let I and I' be two systems of rows, and let J and J' be two systems of columns:

$$I = \{i, i+1, \ldots, i+k\},$$
$$J = \{j, j+1, \ldots, j+l\},$$
$$I' = \{i', i'+1, \ldots, i'+k'\},$$

$$J' = \{j', j'+1, \dots, j'+l'\},$$

where $i + k < j' \leqslant i'$ and $j' + l' \leqslant i' + k' < j$. Suppose the minors $\mathbb{X}_I^{J'}$ and $\mathbb{X}_{I'}^{J}$ are bordered by zeros from below and left in $\mathbb{X}$, where $\mathbb{X}_I^{J'}$ is the minor of the matrix $\mathbb{X}$ with systems of rows I and columns J'. In other words, if for any positive root (a, b) one of the following conditions holds

1) $a \in I$ and $b < j'$,
2) $a \in I'$ and $b < j$,
3) $a > i + k$ and $b \in J'$,
4) $a > i' + k'$ and $b \in J$,

then the root (a, b) is not in the system M. Let $k + k' = l + l'$. The following determinant is one of order $k + k' + 2$:

$$D_{I,I'}^{J,J'} = \left|\begin{array}{c|c} \mathbb{X}_I^{J'} & (\mathbb{X}^2)_I^J \\ \hline 0 & \mathbb{X}_{I'}^{J} \end{array}\right|, \tag{2}$$

where the minor $\mathbb{X}_I^J$ of the matrix $\mathbb{X}$ is formed by systems of rows I and columns J.

Definition 2.1. The determinant $D_{I,I'}^{J,J'}$ is called a *combined minor.*

Lemma 2.2. *A combined minor* (2) *is an N-invariant.*

Proof. A polynomial H is an N-invariant if H is an invariant under the adjoint action of all one-parameter subgroups $g_m(t) = E + tE_{m,m+1}$, $1 \leqslant m < n$, where E is a matrix unit. The adjoint action on the matrix $\mathbb{X}$ by $g_m(t)$ reduces to the composition of two transformations:

1) the row with number $m + 1$ multiplied by t is added to the row m of the matrix $\mathbb{X}$;
2) the column with the number m multiplied by $-t$ is added to the column with the number $m + 1$ of the matrix $\mathbb{X}$.

We show that the polynomial (2) is an invariant under the adjoint action of $g_{m,m+1}(t)$. Without loss of generality, it can be assumed that $i+k = j'-1$, $i' + k' = j - 1$ and the minors $\mathbb{X}_I^{J'}$, $\mathbb{X}_{I'}^{J}$, and $(\mathbb{X}^2)_I^J$ have the following form

$$\mathbb{X}_I^{J'} = \begin{vmatrix} x_{i,j'} & x_{i,j'+1} & \dots & x_{i,j'+l'} \\ x_{i+1,j'} & x_{i+1,j'+1} & \dots & x_{i+1,j'+l'} \\ \dots & \dots & \dots & \dots \\ x_{i+k,j'} & x_{i+k,j'+1} & \dots & x_{i+k,j'+l'} \end{vmatrix},$$

$$\mathbb{X}^{J}_{I'} = \begin{vmatrix} x_{i',j} & x_{i',j+1} & \dots & x_{i',j+l} \\ x_{i'+1,j} & x_{i'+1,j+1} & \dots & x_{i'+1,j+l} \\ \dots & \dots & \dots & \dots \\ x_{i'+k',j} & x_{i'+k',j+1} & \dots & x_{i'+k',j+l} \end{vmatrix},$$

$$(\mathbb{X}^2)^{J}_{I} = \begin{vmatrix} \widetilde{x}_{i,j} & \widetilde{x}_{i,j+1} & \dots & \widetilde{x}_{i,j+l} \\ \widetilde{x}_{i+1,j} & \widetilde{x}_{i+1,j+1} & \dots & \widetilde{x}_{i+1,j+l} \\ \dots & \dots & \dots & \dots \\ \widetilde{x}_{i+k,j} & \widetilde{x}_{i+k,j+1} & \dots & \widetilde{x}_{i+k,j+l} \end{vmatrix},$$

where $\widetilde{x}_{a,b} = \sum\limits_{p=j'}^{i'+k'} x_{a,p}x_{p,b}$ for any $a \in I$ и $b \in J$. If $m < i$ or $m+1 > j+l$, then the adjoint action of the element $g_{m,m+1}(t)$ doesn't change the minor (2).

1. Suppose $i \leqslant m < i+k$; then the action of the element $g_{m,m+1}(t)$ changes the variables $x_{a,b}$ of the minor $D^{J,J'}_{I,I'}$ such that $a = m$ and doesn't change the other variables. Therefore

   $$g_{m,m+1}(t)\, x_{m,p} = x_{m,p} - t x_{m+1,p},\ \ j' \leqslant p \leqslant i' + k';$$

   $$g_{m,m+1}(t) \sum_{p=j'}^{i'+k'} x_{m,p}x_{p,b} = \sum_{p=j'}^{i'+k'} x_{m,p}x_{p,b} - t \sum_{p=j'}^{i'+k'} x_{m+1,p}x_{p,b}.$$

   Then we have

   $$g_{m,m+1}(t)\mathbb{X}^{J}_{I'} = \mathbb{X}^{J}_{I'},$$
   $$g_{m,m+1}(t)\mathbb{X}^{J'}_{I} = \mathbb{X}^{J'}_{I} - tD_1.$$

   The order of D_1 is equal to the order of $\mathbb{X}^{J'}_{I}$. The minor D_1 is formed by the columns J' and D_1 has two identical rows with numbers $m + 1$ and m. Further,

   $$g_{m,m+1}(t)(\mathbb{X}^2)^{J}_{I} = (\mathbb{X}^2)^{J}_{I} - tD_2,$$

   where the order of D_2 is equal to the order of $(\mathbb{X}^2)^{J}_{I}$. The minor D_2 has two identical rows with numbers m and $m + 1$.

   Thus,

   $$g_{m,m+1}(t)D^{J,J'}_{I,I'} = g_{m,m+1}(t) \left|\begin{array}{c|c} \mathbb{X}^{J'}_{I} & (\mathbb{X}^2)^{J}_{I} \\ \hline 0 & \mathbb{X}^{J}_{I'} \end{array}\right| =$$

$$= \left|\begin{array}{c|c} \mathbb{X}_I^{J'} & (\mathbb{X}^2)_I^J \\ \hline 0 & \mathbb{X}_{I'}^J \end{array}\right| - t \left|\begin{array}{c|c} D_1 & D_2 \\ \hline 0 & \mathbb{X}_{I'}^J \end{array}\right|.$$

The minor that consists of D_1, D_2, and $\mathbb{X}_{I'}^J$ has two identical rows; therefore this minor is equal to zero. We have

$$g_{m,m+1}(t) D_{I,I'}^{J,J'} = D_{I,I'}^{J,J'}.$$

2. Let $i + k \leqslant m < j$. For any $a \in I$ and $b \in J$ we have

$$g_{m,m+1}(t) \sum_{p=j'}^{i'+k'} x_{a,p} x_{p,b} = \sum_{p=j'}^{m-1} x_{a,p} x_{p,b} + x_{a,m}(x_{m,b} - t x_{m+1,b}) +$$

$$+ (x_{a,m+1} + t x_{a,m}) x_{m+1,b} + \sum_{p=m+2}^{i'+k'} x_{a,p} x_{p,b} = \sum_{p=j'}^{i'+k'} x_{a,p} x_{p,b},$$

i.e., the adjoint action of the element $g_{m,m+1}(t)$ does not change the minor $(\mathbb{X}^2)_I^J$. Besides, the action of the element $g_{m,m+1}(t)$ adds the column with the number m to the $(m+1)$th column of the minor $\mathbb{X}_I^{J'}$ and the $(m+1)$th row to the mth row of the minor $\mathbb{X}_{I'}^J$. Thus the minor $D_{I,I'}^{J,J'}$ is a $g_{m,m+1}(t)$-invariant.

3. For $j \leqslant m < j + l$, the proof is similarly. □

We shell show that the invariant L_φ is a combined minor (2) for any systems I, I', J, J'.

Definition 2.3. Any root $(i, j) \in M$ satisfying the conditions

$$R_{k-1} < i \leqslant R_k \text{ and } R_k < j \leqslant n,$$

is called a *root lying to the right of the kth block* in $\mathfrak{r}$.

Definition 2.4. Any root $(i, j) \in M$ satisfying the conditions

$$1 \leqslant i \leqslant R_{k-1} \text{ and } R_{k-1} < j \leqslant R_k,$$

is called a *root lying to the above of the kth block* in $\mathfrak{r}$.

Let φ be any root from Φ. The root φ corresponds to an admissible pair (ξ, ξ'). Let $\xi = (i, j)$, $\xi' = (l, m)$, and $i < l$. Assume that φ lies to the right of the kth block in $\mathfrak{r}$, i.e., $R_{k-1} < j \leqslant R_k$. Let the set S_ξ has p roots and let the set $S_{\xi'}$ has q roots. We have $i+p=R_{k-1}$, i.e., the minimal root γ in S_ξ, which has the greatest number of row, lies in the row with number R_{k-1}. Similarly, the root of the base that has the least number of column is also minimal; so we have $l-q=R_k+1$.

Denote

$$I=\{i,i+1,\dots,i+p\},\ J=\{m-q,m-q+1,\dots,m\},$$

$$I'=\{l+1,l+2,\dots,l+q\},\ J'=\{j-p,j-p+1\dots,j-1\}.$$

Then $|I|=p+1,\ |J|=q+1,\ |I'|=q,\ |J'|=p$. Consider the following combined minor

$$\widetilde{L}_\varphi=D^{J,J'}_{I,I'}=\left|\begin{array}{c|c} \mathbb{X}^{J'}_I & (\mathbb{X}^2)^J_I \\ \hline 0 & \mathbb{X}^J_{I'} \end{array}\right|,$$

where $\mathbb{X}^J_I$ is the minor of $\mathbb{X}$ with the systems of rows I and columns J. The order of $\widetilde{L}_\varphi$ is equal to $(p+q+1)$.

Proposition 2.5. *We have the equation* $\widetilde{L}_\varphi=L_\varphi$.

PROOF. By Lemma 2.2, the polynomial $\widetilde{L}_\varphi$ is an invariant. Let

$$p:\ K[\mathfrak{m}]^N\to K[\mathcal{Y}],\quad p(f)=f|_{\mathcal{Y}}$$

be a restriction homomorphism. We calculate the invariants L_φ and $\widetilde{L}_\varphi$ on $\mathcal{Y}$. Let $\mathbb{Y}$ be the formal matrix in which the variables $x_{i,j}$ occupy the positions $(i,j)\in S\cup\Phi$ and the other entries are equal to zero.

Assume that the root φ corresponds to the admissible pair (ξ,ξ'), where

$$\xi=(i,j),\ \xi'=(l,m),$$

and the sets S_ξ, $S_{\xi'}$ consist of the roots

$$S_\xi=\{\xi_1,\xi_2,\dots,\xi_p\},\ S_{\xi'}=\{\xi'_1,\xi'_2,\dots,\xi'_q\},$$

respectively.

Then using elementary transformations of columns, the minor $\mathbb{Y}^{J'}_I$ of the matrix $\mathbb{Y}$ is reduced to the following minor of order $(p+1)\times p$:

$$\left|\begin{array}{cccc} 0 & 0 & \dots & 0 \\ 0 & 0 & \dots & x_{\xi_p} \\ \dots & \dots & \dots & \dots \\ 0 & x_{\xi_2} & \dots & 0 \\ x_{\xi_1} & 0 & \dots & 0 \end{array}\right|.$$

The minor $\mathbb{Y}^{J}_{I'}$ is reduced to the following minor of order $q \times (q+1)$ by elementary transformations of rows:

$$\begin{vmatrix} 0 & 0 & \dots & x_{\xi'_q} & 0 \\ \dots & \dots & \dots & \dots & \dots \\ 0 & x_{\xi'_2} & \dots & 0 & 0 \\ x_{\xi'_1} & 0 & \dots & 0 & 0 \end{vmatrix}.$$

We have

$$\widetilde{L}_\varphi = \pm \left| \begin{array}{cccccccc} 0 & 0 & \dots & 0 & & & & \\ 0 & 0 & \dots & x_{\xi_p} & & & & \\ \dots & \dots & \dots & \dots & & (\mathbb{Y}^2)^J_I & & \\ 0 & x_{\xi_2} & \dots & 0 & & & & \\ x_{\xi_1} & 0 & \dots & 0 & & & & \\ & & & 0 & 0 & \dots & x_{\xi'_q} & 0 \\ & & & \dots & \dots & \dots & \dots & \dots \\ & 0 & & 0 & x_{\xi'_2} & \dots & 0 & 0 \\ & & & x_{\xi'_1} & 0 & \dots & 0 & 0 \end{array} \right| =$$

$$= \pm x_\xi x_\varphi \prod_{s=1}^{p} x_{\xi_s} \cdot \prod_{t=1}^{q} x_{\xi'_t}. \qquad (3)$$

Direct calculations show that the invariant L_φ on $\mathbb{Y}$ is equal to (3).

Let we show that the mapping p is an embedding. Indeed, if $f \in \operatorname{Ker} p$, then $f\,(\mathrm{Ad}_N\,\mathcal{Y}) = 0$. By Theorem 1.9, we have $\mathrm{Ad}_N\,\mathcal{Y}$. Then $f \equiv 0$ and p is an embedding.

The invariant L_φ is equal to $\widetilde{L}_\varphi$ on $\mathcal{Y}$. Therefore L_φ is equal to $\widetilde{L}_\varphi$ on the pre-image $p^{-1}(\mathcal{Y})$. Finally, we get $L_\varphi = \widetilde{L}_\varphi$. □

§3. The additional set of N-invariants

Let $\mathfrak{p}$ be any parabolic subalgebra.

Now we introduce a system of N-invariants that are not in the algebra $K[M_\xi, L_\varphi]_{\substack{\xi\in S\\ \varphi\in\Phi}}$.

Let $(i,j) \in S \cup \Phi$. Denote

$$L_{i,j} = \begin{cases} L_{(i,j)} \text{ if } (i,j) \in \Phi; \\ M_{(a,i)} \cdot M_{(i,j)} \text{ if } (i,j) \in S \text{ and there is a number } a \text{ such that} \\ \hfill (a,i) \in S. \end{cases}$$

Assume that for any numbers $m < l$ and $i < j$ the roots

$$(m, i), (l, i), (m, j), (l, j) \tag{4}$$

are contained in the expanded base. Obviously, the root (m, i) lies in the system Φ. Suppose that the root (m, i) corresponds to some admissible pair $(\xi, \xi') \in Q$, $\xi, \xi' \in S$.

Notation 3.1. Denote

$$A_m^{i,j} = \frac{L_{m+1,i}L_{m,j} - L_{m+1,j}L_{m,i}}{M_\xi}. \tag{5}$$

Now assume that the following condition holds for the roots (4): there is not a number $\widetilde{i}$, $i < \widetilde{i} < j$, such that $(m, \widetilde{i}) \in S \cup \Phi$.

Notation 3.2. Denote

$$B_{m,l}^{i} = \frac{L_{m,j}L_{l,i} - L_{l,j}L_{m,i}}{M_{\xi'}}, \tag{6}$$

$$C_m^i = \frac{L_{m+1,i}L_{m,j} - L_{m+1,j}L_{m,i}}{M_\xi \cdot M_{\xi'}}. \tag{7}$$

Clearly, all rational function (5), (6), (7) are contained in the localization $K[\mathfrak{m}]_{\mathcal{S}}^N$, where the set $\mathcal{S}$ is generated by the minors M_ξ, $\xi \in S$. We shall prove that the functions (5), (6), (7) are polynomials.

Let the root (m, i) be to the right of the kth block in $\mathfrak{r}$, i.e., $R_{k-1} < m \leqslant R_k$. Let the invariant C_m^i is constructed by roots

$$(m, i), (m+1, i), (m, j), (m+1, j),$$

which contained in the expanded base. Then there exists no number $\widetilde{i}$ such that $i < \widetilde{i} < j$ and $(m, \widetilde{i}) \in \Phi$. If the root $(m+1, j)$ lies in the system Φ, then $(m+1, j)$ corresponds to some admissible pair (ξ, ξ'). If $(m+1, j) \in S$, then there exists a root ξ in the column with number $m+1$. Indeed, $(m+1, i) \in \Phi$ corresponds to an admissible pair that contains ξ. In any case, we have that there exist the roots $\xi = (a, m+1)$ and $\xi' = (b, j)$ for some numbers a and b. Consider the sets S_ξ and $S_{\xi'}$. Suppose S_ξ contains p roots and $S_{\xi'}$ contains q roots. Since the root ξ lies to the above of the kth block in $\mathfrak{r}$, then we have $a + p = R_{k-1}$, i.e., the root from S_ξ that has the greatest number of row is contained in the row R_{k-1}. Similarly, $j - q = R_k + 1$.

Example 3.3. Suppose the reductive subalgebra $\mathfrak{r}$ is formed by the blocks $(3, 1, 4, 1, 2, 3)$. We have the following diagram.



Diagram 2

Consider the invariant C_5^{11}. The invariant corresponds to the roots

$$(5,11),(6,11),(5,14),(6,14)$$

from the expanded base. By the above notations, we have $\xi=(2,6)$ and $\xi'=(6,14)$, $p=2$, and $q=5$.

Denote

$$I=\{a,a+1,\dots,a+p\},\ J=\{j-q,j-q+1,\dots,j\},$$

$$I'=\{b+2,b+3,\dots,b+q\},\ J'=\{m-p+1,m-p+2\dots,m-1\}.$$

Then $|I|=p+1,\ |J|=q+1,\ |I'|=q-1,\ |J'|=p-1$. Consider the following combined minor of order $(p+q)$:

$$\widetilde{C}_m^{\,i}=D_{I,I'}^{J,J'}=\left|\begin{array}{c|c}\mathbb{X}_I^{J'} & (\mathbb{X}^2)_I^J\\ \hline 0 & \mathbb{X}_{I'}^J\end{array}\right|,$$

where the minor $\mathbb{X}_I^J$ of the formal matrix is formed by the intersection of rows I and columns J.

Example 3.4. The polynomial $\widetilde{C}_5^{11}$ of Example 3.3 is formed in the following way.

$$\left|\begin{array}{c|c}\mathbb{X}_{2,3,4}^{4} & (\mathbb{X}^2)_{2,3,4}^{9,10,11,12,13,14}\\ \hline 0 & \mathbb{X}_{8,9,10,11}^{9,10,11,12,13,14}\end{array}\right|.$$

The reader will easily prove that $\widetilde{C}_5^{11} = C_5^{11}$.

Proposition 3.5. *We have* $C_m^i = \widetilde{C}_m^i$ *for any numbers* m *and* i.

PROOF. Let us show that $\widetilde{C}_m^i = C_m^i$.

Let

$$p:\ K[\mathfrak{m}]^N \to K[\mathcal{Y}], \quad p(f) = f|_{\mathcal{Y}}$$

be the the restriction homomorphism. We show that p is an embedding. Indeed, if $f \in \operatorname{Ker} p$, then $f\,(\operatorname{Ad}_N \mathcal{Y}) = 0$. Since by Theorem 1.9, $\operatorname{Ad}_N \mathcal{Y}$ is contained a Zariski-open subset, then $f \equiv 0$. Therefore p is an embedding.

By Lemma 2.2, the minor $\widetilde{C}_m^i$ is an invariant. We show that the invariant L_φ is equal to $\widetilde{L}_\varphi$ on $\mathcal{Y}$. Let $\mathbb{Y}$ be the formal matrix such that the entries (i,j) in $\mathbb{Y}$ contain the variables $x_{i,j}$ if $(i,j) \in S \cup \Phi$ and the other entries is filled by zeros. We calculate the polynomial $\widetilde{C}_m^i$ and the invariant C_m^i on $\mathbb{Y}$.

Now assume that the roots

$$\varphi_1 = (m,i),\ \varphi_2 = (m+1,i),\ \varphi_3 = (m,j),\ \varphi_4 = (m+1,j)$$

are in the expanded base and there exists no number $\widetilde{i}$ such that $i < \widetilde{i} < j$ and $(m,\widetilde{i}) \in \Phi$. Suppose the root $\varphi = (m+1,j)$ corresponds to the admissible pair

$$\xi = (a, m+1),\ \xi' = (b,j).$$

Assume that the sets S_ξ and $S_{\xi'}$ consist of the roots

$$S_\xi = \{\xi_1, \xi_2, \ldots, \xi_p\},\ S_{\xi'} = \{\xi'_1, \xi'_2, \ldots, \xi'_q\},$$

where the roots ξ_p and ξ'_q lie in the mth column and in the $(b+1)$th row, respectively. Then by elementary transformations of columns, the minor $\mathbb{Y}_I^{J'}$ of the matrix $\mathbb{Y}$ is reduced to the following minor of order $(p+1) \times (p-1)$:

$$\begin{vmatrix} 0 & 0 & \ldots & 0 \\ 0 & 0 & \ldots & 0 \\ 0 & 0 & \ldots & x_{\xi_{p-1}} \\ \ldots & \ldots & \ldots & \ldots \\ 0 & x_{\xi_2} & \ldots & 0 \\ x_{\xi_1} & 0 & \ldots & 0 \end{vmatrix},$$

By elementary transformations of rows, the minor $\mathbb{Y}_{I'}^{J}$ is reduced to the following minor of order $(q-1) \times (q+1)$:

$$\begin{vmatrix} 0 & 0 & \ldots & x_{\xi'_{q-1}} & 0 & 0 \\ \ldots & \ldots & \ldots & \ldots & \ldots & \ldots \\ 0 & x_{\xi'_2} & \ldots & 0 & 0 & 0 \\ x_{\xi'_1} & 0 & \ldots & 0 & 0 & 0 \end{vmatrix}.$$

We have

$$\widetilde{C}^i_m = \pm \left| \begin{array}{cccccccccc} 0 & 0 & \ldots & 0 & & & & & & \\ 0 & 0 & \ldots & 0 & & & & & & \\ 0 & 0 & \ldots & x_{\xi_{p-1}} & & & (\mathbb{Y}^2)^J_I & & & \\ \ldots & \ldots & \ldots & \ldots & & & & & & \\ 0 & x_{\xi_2} & \ldots & 0 & & & & & & \\ x_{\xi_1} & 0 & \ldots & 0 & & & & & & \\ & & & & 0 & 0 & \ldots & x_{\xi'_{q-1}} & 0 & 0 \\ & & & & \ldots & \ldots & \ldots & \ldots & \ldots & \ldots \\ & 0 & & & 0 & x_{\xi'_2} & \ldots & 0 & 0 & 0 \\ & & & & x_{\xi'_1} & 0 & \ldots & 0 & 0 & 0 \end{array} \right| =$$

$$= \pm \prod_{s=1}^{p-1} x_{\xi_s} \cdot \prod_{t=1}^{q-1} x_{\xi'_t} \cdot \left| \begin{array}{cc} x_\xi x_{\varphi_2} & x_\xi x_{\varphi_4} \\ x_{\xi_p} x_{\varphi_1} & x_{\xi_p} x_{\varphi_3} \end{array} \right| =$$

$$= \pm x_\xi \prod_{s=1}^{p} x_{\xi_s} \cdot \prod_{t=1}^{q-1} x_{\xi'_t} \cdot (x_{\varphi_2} x_{\varphi_3} - x_{\varphi_1} x_{\varphi_4}). \qquad (8)$$

By direct calculations, C^i_m coincides with (8) on $\mathbb{Y}$. Thus C^i_m coincides with (8) on $\mathcal{Y}$, and we have $\widetilde{C}^i_m$ is equal to C^i_m on the pre-image $p^{-1}(\mathcal{Y})$. So, $C^i_m = \widetilde{C}^i_m$. $\square$

Thus the invariant C^i_m is a polynomial.

Proposition 3.6. *We have* $A^{i,j}_m$, $B^i_{m,l} \in K[\mathfrak{m}]^N$.

Proof.Let us prove that $A^{i,j}_m$ is a polynomial. The proof for $B^i_{m,l}$ is similarly.

Suppose the roots (m,i), (m,j), $(m+1,i)$, and $(m+1,j)$, $i<j$, are contained in the expanded base and these roots lie to the right of the kth block in the subalgebra $\mathfrak{r}$. We show that $A^{i,j}_m$ is a polynomial. The proof is by induction.

Assume that there are p roots to the above of the kth block in S. It is easily shown that these roots are the following ones:

$$(i_1, R_{k-1}+1), (i_2, R_{k-1}+2), \ldots, (i_p, R_{k-1}+p)$$

for some $i_1 > i_2 > \ldots > i_p$. Note that we have $i_1 = R_{k-1}$ for the root $(i_1, R_{k-1}+1)$. Suppose there are q roots in S to right of the kth block:

$$(R_k, j_1), (R_k - 1, j_2), \ldots, (R_k - q + 1, j_q)$$

for some $j_1 < j_2 < \ldots < j_q$. We have $j_1 = R_k + 1$. There exist the numbers v and w such that $i = j_v$ and $j = j_w$. Assume that $w - v > 2$. Suppose the root

(m, j_w) corresponds to the admissible pair (ξ, ξ'). We fix the number m. We have the following equation

$$L_{m+1,j_v} M_{\xi'} C_m^{j_{w-1}} = A_m^{j_v, j_w} L_{m+1, j_{w-1}} - A_m^{j_v, j_{w-1}} L_{m+1, j_w} \tag{9}$$

for any $1 < w \leqslant q$. By Proposition 3.5, the left part of the equation (9) lies in $K[\mathfrak{m}]$. But $A_m^{j_v, j_w}$ is contained in the ring $K[\mathfrak{m}]$ provided $A_m^{j_v, j_{w-1}} \in K[\mathfrak{m}]$. Finally from Proposition 3.5 the function

$$A_m^{j_v, j_{v+1}} = C_m^{j_v} \cdot M_{\xi'}$$

lies in the ring $K[\mathfrak{m}]$. $\square$

Conjecture 3.7. Suppose the reductive subalgebra $\mathfrak{r}$ consists of three blocks; then the algebra of invariants $K[\mathfrak{m}]^N$ is generated by the polynomials M_ξ, $\xi \in S$, L_φ, $\varphi \in \Phi$, and the elements (5), (6), (7).

§4. The algebra of invariants in the case $(2, k, 2)$

Assume that the reductive subalgebra of the parabolic subalgebra consists of three blocks with sizes $(2, k, 2)$, $k > 3$. We shall prove Conjecture 3.7 in this case.

The roots

$$S = \Big\{(1,4), (2,3), (k+1, k+4), (k+2, k+3)\Big\},$$

$$\Phi = \Big\{(3, k+3), (3, k+4), (4, k+3), (4, k+4)\Big\}$$

form the base and the system Φ, respectively. We have the following diagram for the parabolic subalgebra in the case $(2, k, 2)$.



Diagram 3

Denote

$$M_1 = M_{(2,3)},\ M_2 = M_{(1,4)},\ N_1 = M_{(k+2,k+3)},\ N_2 = M_{(k+1,k+4)};$$

$$L_{11} = L_{\big((2,3),(k+2,k+3)\big)},\ L_{12} = L_{\big((2,3),(k+1,k+4)\big)},$$
$$L_{21} = L_{\big((1,4),(k+2,k+3)\big)},\ L_{22} = L_{\big((1,4),(k+1,k+4)\big)}.$$

The nonzero polynomials (5), (6) are not defined; the invariant (7) is the following one:

$$D = C_3^{k+3} = \frac{L_{12}L_{21} - L_{11}L_{22}}{M_1N_1}. \tag{10}$$

We show that the invariants M_i, N_j, and L_{ij}, $i,j = 1,2$, do not generate the algebra of invariants $K[\mathfrak{m}]^N$. Denote

$$\mathcal{B}_0 = K[M_1, M_2, N_1, N_2, L_{11}, L_{12}, L_{21}, L_{22}].$$

Assume the converse. Then $K[\mathfrak{m}]^N = \mathcal{B}_0$. Therefore from Theorem 1.10 it follows that there exists a polynomial f in 8 variables and there are numbers l_i, $i = 1, \ldots, 4$, such that

$$D = \frac{f(M_1, M_2, N_1, N_2, L_{11}, L_{12}, L_{21}, L_{22})}{M_1^{l_1} M_2^{l_2} N_1^{l_3} N_2^{l_4}}.$$

Using (10), we get the identity, which contradicts the algebraic independence of the polynomials M_i, N_j, and L_{ij} (see Theorem 1.7).

Theorem 4.1. *For any parabolic subalgebra $\mathfrak{p}$ such that the reductive subalgebra of $\mathfrak{p}$ consists of three block with sizes $(2,k,2)$, $k > 3$, it follows that the algebra of invariants $K[\mathfrak{m}]^N$ is generated by the polynomials M_i, N_i, L_{ij}, and D, where $i,j = 1,2$.*

Proof. Denote the set

$$\mathcal{B} = K[M_i, N_j, L_{ij}, D]_{i,j=1,2} \subset K[\mathfrak{m}]^N.$$

We show that $K[\mathfrak{m}]^N \subset \mathcal{B}$.

As above, let $\mathcal{S}$ be the set of denominators generated by the minors M_i, N_i, where $i = 1,2$. From Theorem 1.10 it follows that the localization $K[\mathfrak{m}]^N_{\mathcal{S}}$ coincides with the algebra of Laurent polynomials

$$K\left[(M_i)^{\pm 1},\ (N_i)^{\pm 1},\ L_{ij}\right]_{i,j=1,2}.$$

If $f \in K[\mathfrak{m}]^N$, then there exist natural numbers k_1, k_2, l_1, l_2 such that

$$f \cdot M_1^{k_1} M_2^{k_2} N_1^{l_1} N_2^{l_2} \in \mathcal{B}_0.$$

Suppose for any minor M from the collection $\{M_i, N_i\}_{i=1,2}$ we have that

$$\text{if } f \in K[\mathfrak{m}] \text{ and } M \cdot f \in \mathcal{B}, \text{ then } f \in \mathcal{B};$$

then $f \in \mathcal{B}$.

Let us prove that if $M \cdot f \in \mathcal{B}$, then $f \in \mathcal{B}$ in the case $M = M_1$. The proof for the other cases is similarly. Suppose $f \in K[\mathfrak{m}]^N$ and $M_1 f \in \mathcal{B}$. The polynomial $M_1 f$ is equal to zero on $\operatorname{Ann} M_1$. Let a_i, b_i, c_{ij}, $i, j = 1, 2$, be any numbers in K. We construct a matrix

$$Q_1 = \begin{pmatrix} 0 & 0 & a_2 & 0 & 0 & \dots & 0 & 0 & 0 \\ 0 & 0 & 0 & a_1 & 0 & \dots & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & \dots & 0 & c_{11} & c_{12} \\ 0 & 0 & 0 & 0 & 0 & \dots & 0 & c_{21} & c_{22} \\ 0 & 0 & 0 & 0 & 0 & \dots & 0 & 0 & 0 \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ 0 & 0 & 0 & 0 & 0 & \dots & 0 & 0 & b_2 \\ 0 & 0 & 0 & 0 & 0 & \dots & 0 & b_1 & 0 \\ 0 & 0 & 0 & 0 & 0 & \dots & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & \dots & 0 & 0 & 0 \end{pmatrix}.$$

By direct calculation, we have

$$M_1(Q_1) = 0, \ M_2(Q_1) = a_1 a_2, \ N_1(Q_1) = b_1, \ N_2(Q_1) = -b_1 b_2;$$

$$L_{11}(Q_1) = a_1 c_{21}, \ L_{12}(Q_1) = -a_1 b_1 c_{22},$$
$$L_{21}(Q_1) = a_1 a_2 c_{21}, \ L_{22}(Q_1) = a_1 a_2 b_1 c_{22},$$
$$D(Q_1) = a_1 a_2 (c_{12} c_{21} - c_{11} c_{22}).$$

Consider an algebra of polynomials

$$\mathcal{A} = K[u_1, u_2, v_1, v_2, w_{11}, w_{12}, w_{21}, w_{22}, z]$$

in 9 variables. Since the polynomial $f M_1$ is contained in $\mathcal{B}$, we have there exists a polynomial $p(u_1, u_2, v_1, v_2, w_{11}, w_{12}, w_{21}, w_{22}, z)$ in the algebra $\mathcal{A}$ such that

$$f M_1 = p(M_1, M_2, N_1, N_2, L_{11}, L_{12}, L_{21}, L_{22}, D). \tag{11}$$

Since $f M_1 = 0$ on $\operatorname{Ann} M_1$, we obtain $(f M_1)(Q_1) = 0$. It now follows that

$$0 = (f M_1)(Q_1) = p\Big(0, a_1 a_2, b_1, -b_1 b_2,$$
$$a_2 c_{21}, -a_2 b_1 c_{22}, a_1 a_2 c_{21}, -a_1 a_2 b_1 c_{22}, a_1 a_2 (c_{12} c_{21} - c_{11} c_{22})\Big).$$

By $\mathcal{Z}$ denote a subset in K^9 that consists of the following collections

$$\Big(0, a_1a_2, b_1, -b_1b_2, a_2c_{21}, -a_2b_1c_{22}, a_1a_2c_{21},\\ -a_1a_2b_1c_{22}, a_1a_2(c_{12}c_{21}-c_{11}c_{22})\Big), \text{ where } a_i, b_j, c_{ij} \in K.$$

It can easily be checked that the polynomials u_1 and $w_{12}w_{21}-w_{11}w_{22}$ are equal to zero on $\mathcal{Z}$. We show that the ideal $\mathcal{I}_{\mathcal{Z}} = \Big\{\varphi \in \mathcal{A} : \varphi(\mathcal{Z}) = 0\Big\}$ is generated by u_1 and $w_{12}w_{21}-w_{11}w_{22}$.

Let the ideal $\mathcal{I}$ be generated by the polynomials $u_1, w_{12}w_{21}-w_{11}w_{22}$. Note that $\mathcal{I}$ is a relevant prime ideal. Indeed, the polynomial $w_{12}w_{21}-w_{11}w_{22}$ is a irreducible one. Therefore the algebra

$$\begin{aligned}\mathcal{A}/\mathcal{I} = \mathcal{A}/\langle u_1, w_{12}w_{21}-w_{11}w_{22}\rangle =\\ = K[u_2, v_1, v_2, w_{11}, w_{12}, w_{21}, w_{22}, z]/\langle w_{12}w_{21}-w_{11}w_{22}\rangle\end{aligned} \tag{12}$$

is a domain of integrity. Hence the ideal $\mathcal{I}$ is a relevant prime one.

Since the polynomials $u_1, w_{12}w_{21}-w_{11}w_{22}$ vanish on the set $\mathcal{Z}$, it follows that $\mathcal{I} \subset \mathcal{I}_{\mathcal{Z}}$ and $\operatorname{Ann}\mathcal{I} \supset \mathcal{Z}$. The dimension of the variety $\operatorname{Ann}\mathcal{I}$ is equal to the transcendence degree of the quotient field of the algebra (12) over K, i.e., $\dim \operatorname{Ann}\mathcal{I} = 7$.

Evidently, $\dim \mathcal{Z} = 7$, then $\dim \operatorname{Ann}\mathcal{I} = \dim \mathcal{Z}$. Since $\operatorname{Ann}\mathcal{I} \supset \mathcal{Z}$, we have $\operatorname{Ann}\mathcal{I} = \overline{\mathcal{Z}}$. We show that $\mathcal{I} = \mathcal{I}_{\mathcal{Z}}$. Suppose $g \in \mathcal{I}_{\mathcal{Z}}$. By the Hilbert's theorem on zeros, there exists a natural number N such that $g^N \in \mathcal{I}$. Since $\mathcal{I}$ is a relevant prime ideal, we have $g \in \mathcal{I}$. Hence, $\mathcal{I}_{\mathcal{Z}} \subset \mathcal{I}$. Using $\mathcal{I} \subset \mathcal{I}_{\mathcal{Z}}$, we obtain $\mathcal{I}_{\mathcal{Z}} = \mathcal{I}$.

Thus since $p|_{\mathcal{Z}} = 0$, it follows that $p \in \mathcal{I}$. Therefore there exist polynomials p_1 and p_2 in the algebra $\mathcal{A}$ such that

$$p = p_1u_1 + p_2(w_{12}w_{21}-w_{11}w_{22}).$$

Combining this with (11), we obtain

$$fM_1 = p_1M_1 + p_2\Big(L_{12}L_{21} - L_{11}L_{22}\Big) = p_1M_1 + p_2M_1N_1D.$$

Further, we have $f = p_1 + p_2N_1D$. So, $f \in \mathcal{B}$. □

Corrolary 4.2. *The algebra of invariants $K[\mathfrak{m}]^N$ is isomorphic onto the factor algebra*

$$K[X_1, X_2, X_3, X_4, Y_1, Y_2, Y_3, Y_4, Z]/(X_2X_4Z - Y_2Y_3 + Y_1Y_4)$$

in the case $(2, k, 2)$, $k > 3$.

§5. The algebra of invariants in the case $(1, 2, 2, 1)$

Suppose the reductive subalgebra $\mathfrak{r}$ of $\mathfrak{p}$ consists of the blocks with sizes $(1, 2, 2, 1)$. We describe the algebra of invariants in this case. The expanded base consists of the following roots:

$$S = \{(1, 2), (3, 4), (2, 5), (5, 6)\}, \quad \Phi = \{(2, 4), (4, 6)\}.$$

The diagram has the form



Diagram 4

The roots of the expanded base correspond to the following invariants.

$$\begin{gathered}
M_1 = M_{(1,2)} = x_{1,2}, \ M_2 = M_{(3,4)} = x_{3,4}, \\
M_3 = M_{(2,5)} = \begin{vmatrix} x_{2,4} & x_{2,5} \\ x_{3,4} & x_{3,5} \end{vmatrix}, \ M_4 = M_{(5,6)} = x_{5,6}; \\
L_1 = L_{(2,4)} = x_{1,2}x_{2,4} + x_{1,3}x_{3,4}, \\
L_2 = L_{(4,6)} = x_{3,4}x_{4,6} + x_{3,5}x_{5,6}.
\end{gathered} \tag{13}$$

Consider a polynomial

$$D = x_{1,2}x_{2,4}x_{4,6} + x_{1,2}x_{2,5}x_{5,6} + x_{1,3}x_{3,4}x_{4,6} + x_{1,3}x_{3,5}x_{5,6}.$$

Lemma 5.1. *The polynomial D is an N-invariant, D is equal to the entry $(1, 6)$ of the matrix $\mathbb{X}^3$, and we have*

$$M_2 D = L_1 L_2 - M_1 M_3 M_4. \tag{14}$$

Proof. This lemma can be proved by direct calculations. □

Note that the algebra of invariants $K[\mathfrak{m}]^N$ does not coincide with the algebra

$$\mathcal{B}_0 = K[M_1, M_2, M_3, M_4, L_1, L_2].$$

Indeed, if $K[\mathfrak{m}]^N = \mathcal{B}_0$, then by Theorem 1.10, there exists a polynomial $f(t_1, \ldots, t_6)$ and there are natural numbers l_i, $i = 1, \ldots, 4$, such that

$$D = \frac{f\Big(M_1, M_2, M_3, M_4, L_1, L_2\Big)}{M_1^{l_1} M_2^{l_2} M_3^{l_3} M_4^{l_4}}.$$

Hence using the equation (14), we obtain that the invariants (13) are algebraically depended. This contradicts Theorem 1.7.

Theorem 5.2. *Let the reductive subalgebra consists of the blocks* $(1,2,2,1)$. *Then the algebra of invariants is generated by the invariants* (13) *and by the polynomial* D.

The proof is similarly to the proof of Theorem 4.1.

Corollary 5.3. *The algebra of invariants* $K[\mathfrak{m}]^N$ *is isomorphic onto the factor algebra*

$$K[X_1,X_2,X_3,X_4,Y_1,Y_2,Z]/(X_2Z-Y_1Y_2+X_1X_3X_4)$$

in the case $(1,2,2,1)$.

So, we have proved that the invariant D is the minor of order 1 of the matrix $\mathbb{X}^3$ in the case $(1,2,2,1)$ (see Lemma 5.1). By Proposition 3.5, the invariants C_m^i can be determined as combined minors, which are constructed by the formal matrices $\mathbb{X}$ and $\mathbb{X}^2$. Similarly, the basic invariants M_ξ, $\xi\in S$, and L_φ, $\varphi\in\Phi$, are combined minors (see Proposition 2.5). Conjecture 5.4 generalizes all cases.

Let $\mathfrak{p}$ be any parabolic subalgebra. Let $I_1,I_2,\ldots,I_k$ and $J_1,J_2,\ldots,J_k$ be any collections of rows and columns, respectively, such that we have the following properties.

1. $|I_1|+|I_2|+\ldots+|I_k|=|J_1|+|J_2|+\ldots+|J_k|$.
2. Let I be a set in the collection $\{I_1,I_2,\ldots,I_k,J_1,J_2,\ldots,J_k\}$. If $\min I< i<\max I$, then $i\in I$.
3. For any $l=1,\ldots,k-1$ we have
$$\max I_l<\min I_{l+1},\ \min J_l>\max J_{l+1}.$$
4. Suppose for a positive root (a,b) and a number l such that $1\leqslant l\leqslant k$, we have one of the following conditions:
   a) $a>\max I_l$ and $b\in J_{k-l+1}$,
   b) $a\in I_l$ and $b<\min J_{k-l+1}$;

   then the root (a,b) is not contained in M. In other words, the minor $\mathbb{X}_{I_i}^{J_{k-i+1}}$ is bordered of zeros to the right and to the below in the matrix $\mathbb{X}$ for any $i=1,\ldots,k$.

We form a determinant, which is defined by the minors of the formal matrix $\mathbb{X}, \mathbb{X}^2, \ldots, \mathbb{X}^k$:

$$D_{I_1,I_2,\ldots,I_k}^{J_1,J_2,\ldots,J_k} = \begin{vmatrix} \mathbb{X}_{I_1}^{J_k} & (\mathbb{X}^2)_{I_1}^{J_{k-1}} & \ldots & (\mathbb{X}^k)_{I_1}^{J_1} \\ 0 & \mathbb{X}_{I_2}^{J_{k-1}} & \ldots & (\mathbb{X}^{k-1})_{I_2}^{J_1} \\ \ldots & \ldots & \ldots & \ldots \\ 0 & 0 & \ldots & \mathbb{X}_{I_k}^{J_1} \end{vmatrix}.$$

Conjecture 5.4. The algebra of invariants $K[\mathfrak{m}]^N$ is generated by the combined minors $D_{I_1,I_2,\ldots,I_k}^{J_1,J_2,\ldots,J_k}$, where the conditions 1–4 hold for $I_1, I_2, \ldots, I_k$ and $J_1, J_2, \ldots, J_k$

References

[B] M. Brion, Representations exceptionnelles des groups semi-simple, *Ann. Scient. Ec. Norm. Sup.* **18** (1985), pp. 345–387.

[GG] M. Goto and F. Grosshans, Semisimple Lie algebras, Lect. Notes in Pure Appl. Math., vol. 38 (1978).

[K] H. Kraft, Geometrische Methoden in der Invariantentheorie, Friedr. Vieweg and Sohn, Braunschweig/Wiesbaden (1985).

[PS] A. N. Panov and V. V. Sevostyanova, Regular N-orbits in the nilradical of a parabolic subalgebra, *Vestnik SamGU*, **7**(57) (2007), pp. 152–161. See also http://arxiv.org/abs/1203.2754.

[PV] V. L. Popov and E. B. Vinberg, Invariant theory, in: *Progressin Science and Technology*, VINITI, Moscow (1989), pp. 137–309.

[R] R. W. Richardson, Conjugacy classes in parabolic subgroups of semisimple algebraic groups, *Bull. London Math. Soc.* **6** (1974), pp. 21–24.

[S] V. V. Sevostaynova, The field of invariants of the adjoint action of the unitriangular group in the nilradical of a parabolic subalgebra. *Zapiski nauchn. seminarov POMI*, vol. 375, 2010, pp. 167–194. (English translation: *Journal of Math. Sciences*, vol. 171, No. 3, 2010, pp. 400–415.) See also http://arxiv.org/abs/1203.3000.

Department of Mechanics and Mathematics, Samara State University, Russia
E-mail address: `victoria.sevostyanova@gmail.com`